Recombinant Mouse CD30 Ligand/TNFSF8 Protein AssayGeni (Fc Tag)



RPES8462

Product Information

Product SKU: RPES8462 Expression Host: Mammalian Size: 20μg

Tag: C-Fc Reactivity: Mouse Accession: P32972

Additional Information

Calculated MW: 43.8 kDa Observed MW: 55-65 kDa

Sequence: Gln68-Asp239

Protein Information

Background: CD30 ligand (CD30L), also known as CD153 and TNFSF8, is a membrane-associated

glycoprotein belonging to the TNF superfamily and TNFR superfamily , and is a

specific ligand for CD30/TNFRSF8 originally described as a cell surface antigen and a

marker for Hodgkin lymphoma and related hematologic malignancies. CD30L is a

type-II membrane glycoprotein expressed on activated T cells , stimulated monocyte-

macrophages , granulocytes , eosinophils , and some Burkitt-like lymphoma cell lines.

CD30L is capable of transducing signals through CD30 on different CD30+ lymphoma

cell lines , and mediates pleiotropic biologic effects including cell proliferation ,

activation , differentiation , as well as cell death by apoptosis. CD30-CD30 ligand

interaction has been suggested to have a pathophysiologic role in malignant

lymphomas , particularly Hodgkin disease , large cell anaplastic lymphomas and

Burkitt lymphomas , and is also involved in activation and functioning of the T cell-

dependent immune response. Thus , CD153 and its receptor CD30 are regarded as

therapeutic targets in hematologic malignancies , autoimmune and inflammatory

diseases.全称

Synonyms: TNFSF, Cd30l, Cd30lg, Tnfsf8, TNFSF8, CD153, CD30 ligand, Tumor necrosis factor

ligand superfamily member 8

Endotoxin: < 1.0 EU/mg of the protein as determined by the LAL method

Formulation: Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Purity: > 90% as determined by reducing SDS-PAGE.

Bio-Activity: Not validated for activity

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.