Recombinant Mouse L-Selectin/CD62L Protein (His AssayGeni



RPES8465

Product Information

Product SKU: RPES8465 Expression Host: Mammalian Size: 20μg

Tag: C-His Reactivity: Mouse Accession: P18337

Additional Information

Calculated MW: 32.2 kDa Observed MW: 55-60 kDa

Sequence: Trp39-Asn332

Protein Information

Background:

L-selectin (SELL), also known as CD62L, is a key adhesion molecule that regulates both the migration of leukocytes at sites of inflammation and the recirculation of lymphocytes between blood and lymphoid tissues. It belongs to the selectin family of proteins, and consisting of a large, highly glycosylated, extracellular domain, a single spanning transmembrane domain and a small cytoplasmic tail. L-selectin is the only selectin expressed on leukocytes and mediates a number of leukocyteendothelial interactions. L-selectin acts as a "homing receptor" for leukocytes to enter secondary lymphoid tissues via high endothelial venules. Ligands present on endothelial cells will bind to leukocyte expressing L-selectin, slowing leukocyte trafficking through the blood, and facilitating entry into a secondary lymphoid organ at that point. L-selectin-mediated lymphocyte recirculation is required for maintaining the appropriate tissue distribution of lymphocyte subpopulations including naïve and effector subsets such as regulatory T cells. In addition, L-selectinmediated entry into peripheral lymph nodes is required for optimal induction of lymphocyte homeostatic proliferation during lymphopenia. Importantly, L-selectin has been shown to have both adhesive and signaling functions during leukocyte migration. L-selectin has also been shown to mediate leukocyte recruitment during chronic inflammatory and autoimmune diseases and thus is a potential therapeutic target for drug development.

Synonyms: Lymphocyte antigen, Leukocyte adhesion molecule, Ly-m, CD62 antigen-like family

member L, CD62L, LAM1, LAM-1, LECAM1, LECAM-1, LEU8, Leu-8, Leukocyte

adhesion molecule 1, LSEL, L-selectin, Lyam1, Lyam-1, Ly-m22, Lymph node homing

receptor, Lymphocyte antigen 22, PLNHR, TQ1, Lnhr, Ly-22, Ly22, Sell, LECAM-1,

LAM-1, Ly-22, MEL-14

Endotoxin: < 1.0 EU/mg of the protein as determined by the LAL method

Formulation: Lyophilized from a 0.2 μm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Purity: > 90% as determined by reducing SDS-PAGE.

Bio-Activity: Not validated for activity

Storage: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at < -20°C for 3 months.