

Product Information

Product SKU:	RPES8470	Expression Host:	Mammalian	Size:	20µg
Tag:	C-Fc	Reactivity:	Mouse	Accession:	O55237

Additional Information

Calculated MW:	41.3 kDa	Observed MW:	45 kDa
Sequence:	Ser45-Ile193		

Protein Information

Background: CD70, a member of the tumor necrosis factor superfamily, is restricted to activated T- and B-lymphocytes and mature dendritic cells. Binding of CD70 to its receptor, CD27, is important in priming, effector functions, differentiation and memory formation of T-cells as well as plasma and memory B-cell generation. Tight control of CD70 expression is required to prevent lethal immunodeficiency. By selective transcription, CD70 is largely confined to activated lymphocytes and dendritic cells (DC). As a type II transmembrane receptor, CD70 is normally expressed on a subset of B, T and NK cells, where it plays a costimulatory role in immune cell activation. Immunohistochemical analysis of CD70 expression in multiple carcinoma types. The restricted expression pattern of CD70 in normal tissues and its widespread expression in various malignancies makes it an attractive target for antibody-based therapeutics. Investigations to exploit CD70 as a cancer target have led to the identification of potential antibody-based clinical candidates.

Synonyms: Tnfsf, Cd27l, Cd27lg, Tnfsf7, Cd70, TNFSF7

Endotoxin: < 1.0 EU/mg of the protein as determined by the LAL method

Formulation: Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.

Purity: > 90% as determined by reducing SDS-PAGE.

Bio-Activity: Not validated for activity

Storage:

Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.