

RPES8492

## Product Information

<b>Product SKU:</b> RPES8492	<b>Expression Host:</b> Mammalian	<b>Size:</b> 20µg
<b>Tag:</b> C-Fc	<b>Reactivity:</b> Mouse	<b>Accession:</b> P13597

## Additional Information

<b>Calculated MW:</b> 75.2 kDa	<b>Observed MW:</b> 90 kDa
<b>Sequence:</b> Gln28-Asn485	

## Protein Information

**Background:** Intercellular adhesion molecule-1 (ICAM-1, or CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and animal models of atherosclerosis. Additionally, ICAM-1 has been implicated in the progression of autoimmune diseases. ICAM-1 is a ligand for LFA-1 (integrin). When activated, leukocytes bind to endothelial cells via ICAM-1/LFA-1 interaction and then transmigrate into tissues. Presence with heavy glycosylation and other structural characteristics, ICAM-1 possesses binding sites for some immune-associated ligands and serves as the binding site for entry of the major group of Human Rhinovirus (HRV) into various cell types. ICAM-1 also becomes known for its affinity for Plasmodium falciparum-infected erythrocytes (PFIE), providing more of a role in infectious disease. Previous studies have shown that ICAM-1 is involved in inflammatory reactions and that a defect in ICAM-1 gene inhibits allergic contact hypersensitivity.

**Synonyms:** MyD, P3., MALA, ICAM-1/CD, MAL, CD54, Icam, Ly-4, Ly-47, Icam-1, MALA-2, BB2, ICAM-1/CD54, Intercellular adhesion molecule 1, MyD10, P3.58, Icam1, Ly-47, MALA-2, CD54, Icam, Icam-1, Ly-4

Contact Details | Dublin, Ireland

Email: [techsupport@assaygenie.com](mailto:techsupport@assaygenie.com) | Web: [www.assaygenie.com](http://www.assaygenie.com)

Copyright © 2024 Assay Genie Ltd, All Rights Reserved. All information / detail is correct at time of going to print.

<b>Endotoxin:</b>	< 1.0 EU/mg of the protein as determined by the LAL method
<b>Formulation:</b>	Lyophilized from a 0.2 µm filtered solution in PBS with 5% Trehalose and 5% Mannitol.
<b>Purity:</b>	> 90% as determined by reducing SDS-PAGE.
<b>Bio-Activity:</b>	Not validated for activity
<b>Storage:</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.