Anti-STAB1 (Bexmarilimab biosimilar) mAb



HDBS0481

| Product Information | | | | |
|---------------------------|--|---|---|--|
| HDBS0481 | Host Species: | Whole mAb | Size: | 100 µg |
| STAB1 | Reactivity : | Human | | |
| mation | | | | |
| Conjugate: Unconjugated U | | niprot: | Q9NY15 | |
| lgG4 | | | | |
| ion | | | | |
| | | | e in therapeutic o | or diagnostic procedures |
| FEX | FEX1;HRFT;FEEL1;FEEL-1;FELE-1;SCARH2;STAB-1;CLEVER-1 | | | |
| e: FP | FP 1305;FP-1305;FP1305 | | | |
| Dilutions: - | | | | |
| - | | | | |
| Res | Research use only | | | |
| Lyo | Lyophilized from sterile PBS, pH 7.4. Normally 5 % – 8% trehalose is added as | | | |
| inst Sto | protectants before lyophilization. Please see Certificate of Analysis for specific instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature. | | | |
| | HDBS0481 STAB1 mation Unconjuga IgG4 ion Res for FEX FP Dilutions: - Res for Pilutions: 500 FEX FP | HDBS0481 Host Species: STAB1 Reactivity: mation Uncorjugated J IgG4 Uncorjugated rom so ranimals Research grade biosin for humans or animals FEX1;HRFT;FEEL1;FEEL FP 1305;FP-1305;FP13 Dilutions: - FEX1;HRFT;FEEL1;FEEL FP 1305;FP-1305;FP13 F0 - F1 - F2 | HDBS0481 Host Species Whole mAb STAB1 Reactivity: Human mation Unconjugated Uniprot: IgG4 Uniprot: IgG4 Iniprot: IgG4 Iniprot: | HDBS0481 Host Species Whole mAb Size: STAB1 Reactivity Human mation Unconjugated Uniprot: Q9NY15 IgG4 Uniprot: Q9NY15 IgG4 Mesearch grade biosimilar. Not for use in therapeutic of for humans or animals. FEX1;HRFT;FEEL1;FEEL-1;FELE-1;SCARH2;STAB-1;CLEVEF FF 1305;FP-1305;FP1305 FEX1;HRFT;FEEL1;FEEL-1;FELE-1;SCARH2;STAB-1;CLEVEF FF 1305;FP-1305;FP1305 FU ions i feconstilition. Kesearch use only Lyophilized from sterile PBS, pH 7.4. Normally 5 % - & protectants before lyophilization. Please see Certificat instructions of reconstitution. Store at -20°C to -80°C for 12 months in lyophilized for |